

LANDAU SINGULARITY AND THE INSTABILITY OF VACUUM STATE IN QED

MOFAZZAL AZAM

*THEORETICAL PHYSICS DIVISION, CENTRAL COMPLEX, BHABHA ATOMIC
RESEARCH CENTRE
TROMBAY, MUMBAI-400085, INDIA*

Abstract

In this paper we argue that the Landau singularity of the coupling constant in QED reflects the instability of the vacuum state, in other words the perturbative ground state of QED. We do not have a rigorous mathematical proof. We rather provide strong arguments derived from the theory of stability of matter developed by Lieb *et al* on the one hand and the theory of weakly interacting fermions developed by Feldman *et al* on the other hand.

PACS numbers:03.65.-w,11.01.-z,12.20.-m

Landau singularity of the coupling constant in Quantum Electrodynamics has been known for more than four decades [1]. The original derivation of the singularity was based on the summation of one loop diagrams of vacuum polarization tensor for photons in the perturbation theory. In the early days the validity of such an expansion was looked upon sceptically by many people including the authors themselves. However, after the advent of $1/N$ expansion technique, it was soon realised that this singularity appears at the leading order in $1/N_f$ expansion where N_f is the number flavour or number of specieses of electrons. This implies that in the infinite flavour limit this singularity is exact. There have been

attempts to interpret this singularity in many ways. Landau and Pomeranchuk tried to argue that this singularity reflects the fact that at short distances strong vacuum polarization effects screen the electric charge completely. Others have called it Landau ghost reflecting the internal inconsistency of Quantum Electrodynamics [2]. In this paper, we shall argue that Landau singularity reflects the instability of vacuum, in other words the perturbative ground state of QED. We do not have a rigorous mathematical proof. We rather rely on 'analogy and precedence', the only other method available to a theoretical physicist in the absence of a mathematical proof. For our 'analogy and precedence' we refer to the theory of stability of matter developed by Lieb *et al* on the one hand and the theory of weakly interacting fermions developed by Feldman *et al* on the other hand.

In quantum theory a many-body system with ground state energy E_0 is called thermodynamically stable or simply stable if E_0/N is bounded below when the number of particles, $N \rightarrow \infty$. In this context, the question of stability of matter consisting of negatively charged electrons and positively charged nuclei is very important. It was in 1967 when Dyson and Lenard proved that, in the framework of nonrelativistic quantum mechanics, matter consisting of N electrons and K static point nuclei is stable [3]. Subsequently, Lieb and his collaborators have made a very detailed investigation of the stability of matter in nonrelativistic as well as relativistic case ([4,5] and references therein). These studies seem to suggest that at high enough energies Quantum Electrodynamics may not be a well behaved theory. Here we argue that there exists one approximation scheme for QED in which it is really so - the perturbative ground state of QED is unstable. This is the large flavour limit of QED, where number of flavours N_f tends to infinity, while $e^2 N_f$ is held fixed, " e " is the QED coupling constant. Thermodynamic stability of a system of particles interacting via electromagnetic interaction is associated with the control of the short distance behaviour of the interaction. In the nonrelativistic limit, zero point kinetic energy of the fermions controls this short distance behaviour - however, in the relativistic limit, there is a need for certain bounds on the value of the fine structure constant [4–6]. In the large flavour limit of QED, Landau singularity becomes exact and it is well known that this is also associated with the

short distance behaviour of the electromagnetic interaction [1] . We refer to the analogy of this model with the theory of weakly interacting electron system analysed by Feldman *et al* using renormalisation group technique. We find an exact analogy between the Landau singularity in QED and the singularity of effective coupling constant in some angular momentum channel in the condensed matter system. In the condensed matter system, we know that the singularity reflects the instability of Landau-Fermi liquid ground (generally called the BCS instability). This analogy clearly suggests that the Landau singularity reflects the instability of ground state in QED. We shall argue in this paper using the example of large flavour QED, that the singularity of effective running coupling constant in the second quantized quantum theory is the equivalent of thermodynamic instability in many-body systems of particles in the first quantized quantum theory.

It should be mentioned here that the stability of ground state and the possible existence of an ultra-violet fixed point has been studied extensively in the lattice formulation of QED in the wide range of values of the fermion flavour N_f by Kogut *et al* [8,9]. On the other hand in massless QED in the continuum, Miransky argues that there exists a chiral symmetry breaking phase [10] . However, in these studies Landau singularity plays no role. There have been some recent studies of Landau singularity using lattice formulation of QED by Gockeler *et al* [11]. These studies seem to suggest that chiral symmetry breaking allows QED to escape Landau singularity. But then chiral symmetry breaking, as their study shows, seems to be intimately connected with triviality of QED. Our approach is motivated by the problems in the theory of stability of matter as the following section would elaborate. We are interested in finding the meaning of Landau singularity rather than finding a way to escape it. (However, we do believe that it is important to investigate the consequences of escaping the singularity as the study in [11] seem to suggest) In our approach Landau singularity is not treated as a ghost as in many other publications but as a signature of instability of the underlying ground state.

I. THE LARGE FLAVOUR QED

The stability of many-body system of fermions corresponding to large flavour QED can be considered in two different ways. One way is to assume that at lower energies only a few flavours are excited. In that case it becomes similar to the single flavour case. In this way we avoid confusion that may arise due to the double limit, number of particles N and number flavour N_f , both tending to infinity. At high energies when we need to consider QED, all specieses of fermions are excited and to avoid additional complication, they can be assumed to be of the same mass. However, we believe this is not the correct way- here we miss the role played by flavour in a many-body theory. It is possible to consider the quantum mechanical many-body problem with large flavours by carefully taking care of the double limit mentioned above. Let us consider a region of linear dimension R where there are N number of fermions with N_f flavours. Assume $N \gg N_f$. Taking into account the the Pauli exclusion principle for the case involoving multflavour fermions in 3-dimensional space, the kinetic energy K can be written [12] as $K \approx \frac{\hbar^2}{2m} \frac{N^{5/3}}{R^2 N_f^{2/3}}$. The potential energy U due to Coulomb interaction is given by [4], $U = -a_0 \frac{e^2 N^{4/3}}{R}$, $a_0 > 0$ is a numerical constant. The total energy, $E(R)$, is given by

$$E(R) = K + U \approx \frac{\hbar^2}{2m} \frac{N^{5/3}}{R^2 N_f^{2/3}} - a_0 \frac{e^2 N^{4/3}}{R} \quad (1)$$

The ground state is obtained by minimizing the total energy, $E(R)$, with respect to R .

$$\begin{aligned} E_0 &\approx -\frac{ma_0^2}{\hbar^2} (e^2 N_f^{1/3})^2 N ; \quad \epsilon_0 = E_0/N \approx -\frac{ma_0^2}{\hbar^2} (e^2 N_f^{1/3})^2 \\ R_0 &\approx \frac{\hbar^2}{ma_0} \left(\frac{N}{N_f}\right)^{1/3} \frac{1}{e^2 N_f^{1/3}} ; \quad r_0 = \frac{R_0}{(N/N_f)^{1/3}} \approx \frac{\hbar^2}{ma_0} \frac{1}{e^2 N_f^{1/3}} \end{aligned} \quad (2)$$

When the number of flavour N_f increases, the size of the small box r_0 decreases and when it approaches Compton wave length $\frac{\hbar}{mc}$, we can no longer use the nonrelativistic quantum mechanics. Therefore, the above estimates are correct only when

$$N_f \ll \frac{1}{a_0^3} \frac{1}{\left(\frac{e^2}{\hbar c}\right)^3} \quad (3)$$

In the relativistic quantum theory of many-body systems, it is almost customary now to take $c|p|$ [4], where c is the velocity of light, as the kinetic energy of individual particles. Therefore, the kinetic energy K for the system of N particles with N_f flavour in a region with linear dimension R is given by, $K = \frac{\hbar c N^{4/3}}{R N_f^{1/3}}$. Therefore, in the relativistic case,

$$E(R) = K + U \approx \frac{\hbar c N^{4/3}}{R N_f^{1/3}} - a_0 \frac{e^2 N^{4/3}}{R} \quad (4)$$

In this case stability of ground state requires $E(R) > 0$. This leads to the condition,

$$N_f < \text{or} = \frac{1}{a_0^3} \frac{1}{(\frac{e^2}{\hbar c})^3} \quad (5)$$

which in the unit $c = \hbar = 1$, becomes $N_f < \text{or} = \frac{1}{(a_0 e^2)^3}$. At this point a few comments are necessary. We considered a theory with large N and large N_f , and it is implicit in our discussion that we are considering, $N \rightarrow \infty$, but we have said nothing about the flavour N_f . We could have also taken $N_f \rightarrow \infty$ but slower than N . We could also assume that in the limit $N_f \rightarrow \infty$ and $e \rightarrow 0$, $e^2 N_f^{1/3} = \text{constant}$, and satisfies the equation Eq.[5] and conclude that in this limit matter is stable. Mathematically it is correct. But this is only a formal result. e does not tend to zero and N_f , never tends to infinity. The mathematical limit, has the physical implication that when e is small and N_f is large, the relevant parameter that sets the stability criteria is $e^2 N_f^{1/3}$. It is the value of $e^2 N_f^{1/3}$ that decides the stability of the system. When $e^2 N_f^{1/3} > \frac{1}{a_0^3}$, in other words, when the number of flavours $N_f > \frac{1}{(a_0 e^2)^3}$, the **many-body system is thermodynamically unstable**.

In Quantum Electrodynamics we find that the relevant parameter is not $e^2 N_f^{1/3}$ but $e^2 N_f$. In large flavour QED with N_f flavours, when we calculate the connected Green's functions using Feynman diagram technique, we find that one loop diagrams are of order $e^2 N_f$ and higher loop diagrams are of lower order in N_f . This becomes more transparent by carrying out expansion in $1/N_f$. The essential idea is to replace $e \rightarrow e/\sqrt{N_f}$, and consider the Lagrangian,

$$\mathcal{L} = \sum_{i=1}^{N_f} \bar{\Psi}^i \left(i \gamma^\mu \partial_\mu + m - e/\sqrt{N_f} \gamma^\mu A_\mu \right) \Psi^i + \frac{1}{4} F_{\mu\nu}^2 \quad (6)$$

where ψ is the Dirac field, and A_μ and $F_{\mu\nu}$ are the potential and the field strength respectively. In the leading (zeroth) order in $1/N_f$, the photon propagator receives radiative corrections only from the single-loop Feynman diagrams. Higher loop contributions are of order one or more in $1/N_f$. The renormalized running (effective) coupling constant at energy, $k^2 = \Lambda^2$, up to the leading order in $1/N_f$, is given by

$$e_{eff}^2(\Lambda^2) = \frac{e^2}{1 - (e^2/3\pi)\ln\Lambda^2/m^2} \quad (7)$$

where we have used, $e^2 = e^2(k^2 = m^2)$, for the physical coupling constant. One can easily infer that the running coupling constant blows up at, $\Lambda^2 = m^2 \exp(3\pi/e^2)$. This is known as **Landau singularity** in QED [1]. It is easy to see that this singularity is associated with the short distance (large momentum) behaviour of the interaction. To understand the meaning of Landau singularity, here we refer to the renormalisation group analysis of weakly interacting Fermi system with short range potential at finite density and zero temperature by Feldman *et al* [7,13]. The iterative renormalization group transformations leads to the same type of singularity in some suitably defined running coupling constant.

II. FELDMAN MODEL OF WEAKLY INTERACTING ELECTRON GAS

We briefly describe here the Feldman Model of weakly interacting electron gas (see reference [13] for details). The model of weakly interacting electron gas studied by Feldman *et al* is a condensed matter Fermi system in thermal equilibrium at some temperature T (for simplicity, assume $T = 0$) and chemical potential μ . On microscopic scale ($\approx 10^{-8}$ cm), it can be described approximately in terms of non-relativistic electrons with short range two body interactions. The thermodynamic quantities such as conductivity depend only on physical properties of the system at mesoscopic length scale ($\approx 10^{-4}$ cm), and therefore, are determined from processes involving momenta of the order of $\frac{k_F}{\lambda}$ around the Fermi surface, where the parameter, $\lambda \gg 1$, should be thought of as a ratio of meso-to-microscopic length scale. This is generally referred to as the scaling limit (large λ , low

frequencies) of the system. The most important observation of Feldman et. al. is that in the scaling limit, systems of non-relativistic (free) electrons in d spatial dimensions behave like a system of multi-flavoured relativistic chiral Dirac fermions in $1 + 1$ dimensions. The number of flavours $N \approx \text{const. } \lambda^{d-1}$. It is possible then to set up a renormalization group improved perturbation theory in $\frac{1}{\lambda}$ around the non-interacting electron gas, where in, the large number of flavours N , play an important role in actual calculations.

A. Free Electron gas and the Multiflavour Relativistic Fermions in $1 + 1$ Dimensions

Let us consider a system of non-relativistic free electrons in d spatial dimensions with the Euclidean action,

$$S_0(\psi^*, \psi) = \sum_{\sigma} \int d^{d+1}x \psi_{\sigma}^*(x) (i\partial_0 - \frac{1}{2m}\Delta - \mu) \psi_{\sigma}(x) \quad (8)$$

The Euclidean free fermion Green's function, $G_{\sigma\sigma'}^0(x-y)$, where σ and σ' are the spin indices, $x = (t, \vec{x})$ and $y = (s, \vec{y})$, t and s are imaginary times, $t > s$, is given by,

$$G_{\sigma\sigma'}^0(x-y) = \langle \psi_{\sigma}^*(x) \psi_{\sigma}(y) \rangle_{\mu} = -\delta_{\sigma\sigma'} \int (dk) \frac{e^{-ik_0(t-s) + i\vec{k}(\vec{x}-\vec{y})}}{ik_0 - (\frac{k^2}{2m} - \mu)} \quad (9)$$

Where we have used $(dk) = \frac{1}{(2\pi)^{(d+1)}} d^{d+1}k$. In the scaling limit, the leading contributions to $G_{\sigma\sigma'}^0(x-y)$ come from modes whose momenta are contained in a shell $S_F^{(\lambda)}$ of thickness $\frac{k_F}{\lambda}$ around the Fermi surface S_F . In order to approximate the Green's function, let us introduce the new variables $\vec{\omega}$, $p_{\parallel}$, p_0 such that $k_F \vec{\omega} \in S_F$, $p_0 = k_0$ and $\vec{k} = (k_F + p_{\parallel})\vec{\omega}$. If $\vec{k} \in S_F^{(\lambda)}$, then $p_{\parallel} \ll k_F$, and we can approximate the integrand of Eq.(4), by dropping $p_{\parallel}^2$ term in the denominator. In other words,

$$G_{\sigma\sigma'}^0(x-y) = \delta_{\sigma\sigma'} \int \frac{d\sigma(\vec{\omega})}{(2\pi)^{d-1}} k_F^{d-1} e^{ik_F \vec{\omega}(\vec{x}-\vec{y})} G_c(t-s, \vec{\omega}(\vec{x}-\vec{y})) \quad (10)$$

where $d\sigma(\vec{\omega})$ is the uniform measure on unit sphere and

$$G_c(t-s, \vec{\omega}(\vec{x}-\vec{y})) = - \int \frac{dp_0}{2\pi} \frac{dp_{\parallel}}{2\pi} \frac{e^{-ik_0(t-s) + ip_{\parallel} \vec{\omega}(\vec{x}-\vec{y})}}{ip_0 - v_F p_{\parallel}} \quad (11)$$

is the Green's function of chiral Dirac fermion in $1 + 1$ dimension. $v_F = k_F/m$ is the Fermi velocity. The $\vec{\omega}$ -integration in Eq.(5) can be further approximated by replacing it with summation over discrete directions $\vec{\omega}_j$ by dividing the shell $S_F^{(\lambda)}$ into N small boxes $B_{\vec{\omega}_j}, j = 1, \dots, N$ of roughly cubical shape. The box, $B_{\vec{\omega}_j}$, is centered at $\vec{\omega}_j \in S_F$ and has an approximate side length $\frac{k_F}{\lambda}$. The number of boxes, $N = \Omega_{d-1} \lambda^{d-1}$, where Ω_{d-1} is the surface volume of unit sphere in d spatial dimensions. The Green's function is, now, given by

$$G_{\sigma\sigma'}^0(x - y) = -\delta_{\sigma\sigma'} \sum_{\vec{\omega}_j} \int \frac{dp_0}{2\pi} \frac{dp_{\parallel}}{2\pi} \frac{p_{\perp}}{2\pi} \frac{e^{-ip_0(t-s) + i\vec{p}(\vec{x}-\vec{y})}}{ip_0 - v_F p_{\parallel}} \quad (12)$$

where $\vec{p} = p_{\parallel} \vec{\omega} + \vec{p}_{\perp}$ is a vector in $B_{\vec{\omega}_j} - k_F \vec{\omega}_j$ and $p_0 \in \mathcal{R}$. Thus in the scaling limit, the behaviour of a d -dimensional non-relativistic free electron gas is described by $N = \Omega_{d-1} \lambda^{d-1}$ flavours of free chiral Dirac fermions in $1 + 1$ dimensional space-time. The propagator $G_c(t - s, \vec{\omega}(\vec{x} - \vec{y}))$ depends on the flavour index $\vec{\omega}$. But the energy of an electron or hole with momenta $\vec{k}$, depends only on $p_{\parallel}$, where $p_{\parallel} = \vec{k} \cdot \vec{\omega} - k_F$ and $\vec{\omega} = \frac{\vec{k}}{|\vec{k}|}$. It is proportional to $p_{\parallel}$, just as for relativistic fermions in $1 + 1$ dimensions.

B. Renormalization Group Flow and the BCS Instability

Large scale behaviour of the weakly interacting system is described by an effective action. To see how the effective action is calculated, let us consider a system with Euclidean action of the form,

$$S(\psi^*, \psi) = S_0(\psi^*, \psi) + S_I(\psi^*, \psi) \quad (13)$$

where $S_0(\psi^*, \psi)$ is the quadratic part given in Eq.(3) and $S_I(\psi^*, \psi)$ is the quartic interaction term. For a weakly interacting electron gas,

$$S_I(\psi^*, \psi) = g_0 k_F^{1-d} \sum_{\sigma, \sigma'} \int d^{d+1}x d^{d+1}y : \psi_{\sigma}^*(x) \psi_{\sigma}(x) v(\vec{x} - \vec{y}) \delta(x_0 - y_0) \psi_{\sigma'}^*(y) \psi_{\sigma'}(y) \quad (14)$$

The factor k_F^{1-d} ensures that g_0 is dimensionless. The two body potential $v(\vec{x})$ is assumed to be smooth and short range, and therefore, its Fourier transform $\hat{v}(\vec{k})$ is also smooth. In

order to calculate the effective actions, we first split the field variable into slow and fast modes, $\hat{\psi} = \hat{\psi}_< + \hat{\psi}_>$, where $\text{Supp } \hat{\psi}_> \subset R \times (R^d \setminus S_F^{(\lambda)})$ (region $>$) and $\text{Supp } \hat{\psi}_< \subset R \times S_F^{(\lambda)}$ (region $<$). We then integrate out the fast modes using the functional integral for fermions,

$$e^{-S_{eff}(\hat{\psi}_<^*, \hat{\psi}_<)} = \frac{1}{\Xi} \int D\hat{\psi}_>^* D\hat{\psi}_> e^{-S(\hat{\psi}_>^*, \hat{\psi}_<^*, \hat{\psi}_>, \hat{\psi}_<)} \quad (15)$$

Now using the linked cluster theorem, we obtain

$$e^{-S_{eff}(\hat{\psi}_<^*, \hat{\psi}_<)} = \exp(-S_{0,<-} < S_I >_{G_>^0} + \frac{1}{2} < S_I; S_I >_{G_>^0} - \frac{1}{3} < S_I; S_I; S_I >_{G_>^0} - \dots) \quad (16)$$

The abbreviations, $< a; b > < a; b; c >$ etc., denote the connected correlators. The subscripts $G_>^0$ indicate that the expectations $< (\cdot) >_{G_>^0}$ are calculated using infrared cut-off free propagators in accordance with the functional measure, $dP(\hat{\psi}_>^*, \hat{\psi}_>) = (1/\Xi) D\hat{\psi}_>^* D\hat{\psi}_> e^{-S_0(\hat{\psi}_>^*, \hat{\psi}_>)}$. The connected correlators can be evaluated using the Feynman diagram technique. Therefore, the effective action can be calculated once we know the amplitudes of connected Feynman diagrams. It is clear that the S_{eff} contains far more interactions than the original quartic interaction S_I . However, for weakly interacting systems the original coupling remains dominant. To carry out the iterative renormalization group scheme of Feldman *et al*, we choose some large scale $\lambda_0 \ll \frac{1}{\sqrt{g_0}}$ and calculate the effective action, S_{eff} perturbatively to leading order in $\frac{1}{\lambda_0}$. The effective action depends on modes corresponding to wave vectors located in the shell, $S_F^{(\lambda)}$, of width $\frac{k_F}{\lambda_0}$ around the Fermi surface. Now, we divide the shell $S_F^{(\lambda)}$ into $N = \text{const. } \lambda^{d-1}$ cubical boxes, $B_{\vec{\omega}_j}$, of approximate side length $\frac{k_F}{\lambda_0}$. Next, we rescale all the momenta so that, instead of belonging to the boxes $B_{\vec{\omega}_j}$ they are contained in boxes $\tilde{B}_{\vec{\omega}_j}$ of side length $\approx k_F$. These two steps are generally known as decimation of degrees of freedom and rescaling. The renormalization group scheme consists of iteration of these two steps.

Assume that the degrees of freedom corresponding to momenta not lying in $S_F^{(\lambda)}$ have been integrated out. Let $\hat{\psi}_\sigma(k), k \in R \times S_F^{(\lambda)}$, denote these modes. The sector fields are defined as

$$\psi_{\vec{\omega},\sigma}(x) = \int_{R \times B_{\vec{\omega}}} (dk) e^{i(k_0 t - (\vec{k} - k_F \vec{\omega}) \vec{x})} \hat{\psi}_{\sigma}(k) \quad (17)$$

It is easy to see that $\psi_{\sigma}(x) = \sum_{\vec{\omega}} e^{ik_F \vec{\omega} \vec{x}} \hat{\psi}_{\vec{\omega},\sigma}(x)$. Inserting the Fourier transform of sector fields in Eq.(3) and Eq.(9), and carrying out some algebraic manipulations, we obtain

$$S_0 = - \sum_{\vec{\omega},\sigma} \int_{\mathcal{R} \times (B_{\vec{\omega}} - \|\mathcal{F}\vec{\omega})} (dp) \hat{\psi}_{\vec{\omega},\sigma}^*(p) (ip_0 - v_F \vec{\omega} \vec{p} + O(\frac{1}{\lambda^2})) \hat{\psi}_{\vec{\omega},\sigma}(p) \quad (18)$$

$$S_I = \frac{g_0}{2} k_F^{1-d} \sum_{\vec{\omega}_1, \dots, \vec{\omega}_4; \sigma, \sigma'} \hat{v}(k_F(\vec{\omega}_1 - \vec{\omega}_4)) \int (dp_1) \dots (dp_4) (2\pi)^{d+1} \delta(p_1 + p_2 - p_3 - p_4) \\ \hat{\psi}_{\vec{\omega}_1, \sigma}^*(p_1) \hat{\psi}_{\vec{\omega}_2, \sigma'}^*(p_2) \hat{\psi}_{\vec{\omega}_3, \sigma'}(p_3) \hat{\psi}_{\vec{\omega}_4, \sigma}(p_4) + \text{terms of higher order in } \frac{1}{\lambda} \quad (19)$$

Using cluster expansions to integrate out the degrees of freedom corresponding to momenta outside the shell $S_F^{(\lambda_0)}$, one can show that, at scale $\frac{k_F}{\lambda}$, the effective action has the form given by Eq.(13) and Eq.(14) except that $\hat{v}(k_F(\vec{\omega}_1 - \vec{\omega}_4))$ is replaced by a coupling function $g(\vec{\omega}_1, \vec{\omega}_2, \vec{\omega}_3, \vec{\omega}_4) \approx \hat{v}(k_F(\vec{\omega}_1 - \vec{\omega}_4))$ with $\vec{\omega}_1 + \vec{\omega}_2 = \vec{\omega}_3 + \vec{\omega}_4$.

Next, one considers the rescaling of the fields and the action. The fields are rescaled in such a way that the supports of the Fourier transformed "sector fields" are boxes, $\tilde{B}_{\vec{\omega}} = \lambda(B_{\vec{\omega}} - k_F \vec{\omega})$, of roughly cubical shape with sides of length k_F , and such that the quadratic part of the action remains unchanged to leading order in $\frac{1}{\lambda}$. The first condition implies that $p \mapsto \tilde{p} = p\lambda$ and $x \mapsto \xi = \frac{x}{\lambda}$. The rescaled sector fields and their Fourier transforms are given by,

$$\tilde{\psi}_{\vec{\omega},\sigma}(\xi) = \lambda^{\alpha} \psi_{\vec{\omega},\sigma}(\lambda \xi) \quad ; \quad \hat{\tilde{\psi}}_{\vec{\omega},\sigma}(\tilde{p}) = \lambda^{\alpha-d-1} \hat{\psi}_{\vec{\omega},\sigma}(\frac{\tilde{p}}{\lambda}) \quad (20)$$

Inserting the scaled Fourier transformed fields into quadratic part of action S_0 , it is easy to see that S_0 remains unchanged if the scaling dimension $\alpha = \frac{d}{2}$. Now, inserting the rescaled fields in the quartic part of the action, we find that the quartic part has scaling dimension $(1-d)$. The quartic terms of higher degree in momenta as well as terms of higher degree in fields appearing in the effective action have smaller scaling dimensions.

Thus the effective action in terms of scaled sector fields is,

$$\begin{aligned}
S_{eff} = & \sum_{\vec{\omega}, \sigma} \int (d\vec{p}) \hat{\psi}_{\vec{\omega}, \sigma}^*(\vec{p}) (i\vec{p}_0 - v_F \vec{\omega} \vec{p}) \hat{\psi}_{\vec{\omega}, \sigma}(\vec{p}) + \frac{1}{2} \frac{1}{\lambda^{d-1}} \sum_{\vec{\omega}_1 + \vec{\omega}_2 = \vec{\omega}_3 + \vec{\omega}_4; \sigma, \sigma'} g(\vec{\omega}_1, \vec{\omega}_2, \vec{\omega}_3, \vec{\omega}_4) \\
& \int (d\vec{p}_1) \dots (d\vec{p}_4) (2\pi)^{d+1} \delta(\vec{p}_1 + \vec{p}_2 - \vec{p}_3 - \vec{p}_4) \hat{\psi}_{\vec{\omega}_1, \sigma}^*(\vec{p}_1) \hat{\psi}_{\vec{\omega}_2, \sigma'}^*(\vec{p}_2) \hat{\psi}_{\vec{\omega}_3, \sigma'}(\vec{p}_3) \hat{\psi}_{\vec{\omega}_4, \sigma}(\vec{p}_4) \\
& + \text{terms of higher order in } \frac{1}{\lambda}
\end{aligned} \tag{21}$$

We see that that the inverse propagator for the sector field is diagonal in $\vec{\omega}$, and it depends only on p_0 and $p_{\parallel} = \vec{\omega} \vec{p}$ but not on $p_{\perp} = \vec{p} - (\vec{\omega} \cdot \vec{p}) \vec{\omega}$.

We are interested in the renormalization group flow equations in the leading order in $\frac{1}{\lambda}$. Therefore, for carrying out the decimation of degrees of freedom, we will be interested only in those diagrams that contribute to the amplitude in the leading order in $\frac{1}{\lambda}$. Before we consider these diagrams, let us consider the possible inter sector scattering geometries. How many independent $g^{(0)}(\vec{\omega}_1, \vec{\omega}_2, \vec{\omega}_3, \vec{\omega}_4)$ exists? For $d = 3$, suppose $\vec{\omega}_3 \neq -\vec{\omega}_4$. On the unit sphere, there are $N^{(0)} = \text{Const.} \lambda_0^{d-1}$ different $\vec{\omega}$'s. But all choices of $\vec{\omega}_1$ and $\vec{\omega}_2$ with $\vec{\omega}_1 + \vec{\omega}_2 = \vec{\omega}_3 + \vec{\omega}_4$ lie on a cone containing $\vec{\omega}_3$ and $\vec{\omega}_4$ with $\vec{\omega}_3 + \vec{\omega}_4$ as the symmetry axis. Therefore, there are $O(\lambda_0^{d-2})$ choices. Only when $\vec{\omega}_3 = -\vec{\omega}_4$ that there are $N^{(0)} = \text{Const.} \lambda_0^{d-1}$ choices. Couplings involving incoming states with $\vec{\omega}_3 \neq -\vec{\omega}_4$ will be represented by $g^{(0)}(\vec{\omega}_1, \vec{\omega}_2, \vec{\omega}_3, \vec{\omega}_4)$. Couplings that involve sectors $\vec{\omega}_3 = -\vec{\omega}_4$ or equivalently $\vec{\omega}_1 = -\vec{\omega}_2$ will be denoted by $g_{BCS}^{(0)}(\vec{\omega}_1, \vec{\omega}_4)$. Because of rotational invariance, $g_{BCS}^{(0)}(\vec{\omega}_1, \vec{\omega}_4)$ is a function of only the angle between $\vec{\omega}_1$ and $\vec{\omega}_4$.

The chemical potential receives corrections from connected diagrams with two external electron lines (self energy correction of the electrons). The contribution of order zero in $\frac{1}{\lambda}$ comes only from the **tadpole** and **turtle** diagrams. These diagrams contain one internal interaction squiggle of order $\frac{1}{\lambda_0^{d-1}}$ and there are $N^{(0)} = \text{Const.} \lambda_0^{d-1}$ choices of the inner particle sector. Therefore, the contribution is of order zero in $\frac{1}{\lambda}$. The amplitude corresponding to these tadpole and turtle diagrams turns out to be p -independent but of order $O(g^{(0)})$. Thus, $\delta\mu_1 = O(g^{(0)}/\lambda_0)$, and the renormalized electron propagator is $G_{\vec{\omega}}^R(p_0, \vec{p}, \vec{\omega}) = -[i\vec{p}_0 - v_F \vec{p} \cdot \vec{\omega} + \lambda_0 \delta\mu_1]^{-1}$.

To find the evolution of the coupling constant $g(\vec{\omega}_1, \vec{\omega}_2, \vec{\omega}_3, \vec{\omega}_4)$, we have to calculate amplitude of diagrams with four external legs. It is found that when $\vec{\omega}_1 + \vec{\omega}_2 = \vec{\omega}_3 + \vec{\omega}_4 \neq 0$, the coupling functions do not flow in the leading order in $\frac{1}{\lambda}$. But for sector indices $\vec{\omega}_1 + \vec{\omega}_2 = \vec{\omega}_3 + \vec{\omega}_4 = 0$, the coupling functions, $g_{BCS}^{(0)}(\vec{\omega}_1, \vec{\omega}_4)$, flow. The diagrams that contribute to the flow equation are the **ladder diagrams** with self energy insertion for the internal electron lines but with no other two legged subdiagram. The amplitude of such a diagram with n interaction squiggles and with zero incoming and outgoing box momenta of the particles is given by,

$$\left(\frac{1}{\lambda^{d-1}}\right)^{n+1} \sum_{\vec{\omega}_1, \dots, \vec{\omega}_n} (-1)^n \beta^n g_{BCS}(\vec{\omega}, \vec{\omega}_n) g_{BCS}(\vec{\omega}_n, \vec{\omega}_{n-1}) \dots g_{BCS}(\vec{\omega}_1, \vec{\omega}')$$

In the equation above, $(\vec{\omega}', -\vec{\omega}')$ and $(\vec{\omega}, -\vec{\omega})$ are sector indices of incoming and outgoing electron lines respectively. Other sector indices correspond to the internal electron lines. β is a strictly positive number coming from the fermion loop integration in the Feynman diagram and is given by $\beta = \int dk_{\perp} dk_{\parallel} dk_0 [k_0^2 + (v_F k_{\parallel} - \lambda \delta \mu_1)^2]^{-1}$. We find that the renormalized value of g_{BCS} is

$$g_{BCS}^{(j+1)}(\vec{\omega}, \vec{\omega}') = g_{BCS}^{(j)}(\vec{\omega}, \vec{\omega}') + \sum_{n=1}^{\infty} \left(\frac{1}{\lambda^{d-1}}\right)^n \sum_{\vec{\omega}_1, \dots, \vec{\omega}_n} (-1)^n \beta_j^n g_{BCS}^{(j)}(\vec{\omega}, \vec{\omega}_n) g_{BCS}^{(j)}(\vec{\omega}_n, \vec{\omega}_{n-1}) \dots g_{BCS}^{(j)}(\vec{\omega}_1, \vec{\omega}') + O\left(\frac{g^{(j)}}{\lambda_j}\right) \quad (22)$$

The explicit expression for flow equation can be obtained by expanding the coupling functions $g_{BCS}(\vec{\omega}, \vec{\omega}') = g_{BCS}(\langle \vec{\omega}, \vec{\omega}' \rangle)$ into spherical harmonics, $g_{BCS}(\langle \vec{\omega}, \vec{\omega}' \rangle) = \sum g_l h_l(\langle \vec{\omega}, \vec{\omega}' \rangle)$. Up to terms of order $\frac{1}{\lambda}$,

$$g_l^{(j+1)} = \frac{g_l^{(j)}}{1 + \beta_j g_l^{(j)}} + O\left(\frac{1}{\lambda}\right) \quad (23)$$

To obtain the differential equation for the R.G. flow, let us define $g_l^{(j)} := g_l^{(\lambda_j)}$, and consider a scale $\lambda = e^t \lambda_0$. Next, define $g_l(t) := g_l^{(e^t \lambda_0)}$. The coefficient $\beta = \beta(t, t')$ vanishes in the limit $t' \searrow t$, therefore, $\beta(t', t) = (t' - t)\gamma(t) + O((t' - t)^2)$. Writing difference equation for the couplings $g_l^{(j+1)}$ and $g_l^{(j)}$, and dividing both sides of the difference equation by $(t' - t)$, we finally obtain

$$\frac{d}{dt}g_l(t) = -\gamma g_l(t)(t)^2 + O(e^{-t}g(t)^2) \quad (24)$$

where $\gamma = \gamma(t) > 0$. It is independent of l and approximately independent of t , and therefore, we set $\gamma = \gamma_0$. The positivity of γ follows from slow monotone growth of $\beta(t' - t)$ in t' . Neglecting the error term, the solution can be written as,

$$g_l(t) = \frac{g_l(0)}{1 + \gamma_0 g_l(0)t}. \quad (25)$$

If there is an angular momentum channel, l , with attractive interactions ($g_l(0) < 0$) the flow diverges at a finite value of the scaling parameter, $t = -(\gamma_0 g_l(0))^{-1}$. This singularity reflects the BCS-instability of the ground state.

III. INSTABILITY OF THE PERTURBATIVE GROUND STATE IN QED

We have seen that in the case of large flavour QED as well as the weakly interacting Fermi system with attractive potential in any one of the angular momentum channel, the running coupling constant blow up at some finite value of their arguments. In the latter case, we know that the perturbative ground state is unstable (BCS instability): perturbation theory is built around the wrong ground state (the non-interacting Fermi gas). The elementary excitations are the electrons and holes, and Fock space is constructed using these single particle states. The BCS instability implies that such excitations do not exist- we require a modification of the excitation spectrum. But this also means that the new stable ground state is not in the eigenstate of particle number operator, $\hat{N}$ and the global $U(1)$ - gauge symmetry associated with the conservation of particle number (excitations) is broken. The BCS ground state of superconductivity [13] contains all these features.

Thus the BCS theory of superconductivity contains two parts :

- 1) The Landau-Fermi liquid ground state is unstable
- 2) There exists another ground state- superconducting state with Cooper pairs as the elementary modes.

In this paper, we have been concerned with the first part of the problem. We have seen that, up to leading order in $1/N$, interactions in the various angular momentum channels decouple, and attractive interaction in any angular momentum channel leads to the instability of (Landau-Fermi liquid) ground state. This analysis in the Feldman theory was carried out using renormalization group technique in the language of "Second Quantization". On the other hand the Lieb theory of stability of matter is discussed in language of "First Quantization". Both the methods address the issue of stability of "Ground State of a Many-Body System". To understand the connection between the two methods, let us consider a realistic Fermi system. The most interesting realistic Fermi system happens to be the liquid He^3 [14]. The minimum of the interatomic potential in He^3 (as well as in He^4) is too close to the atoms and is too shallow and, therefore the zero point is high compared to the potential energy and the helium atoms do not bind to form molecules. However, at higher density (at high pressure) the net potential energy decreases as $-N^2v$ (v being the two body potential) and overcomes the zero point energy which increases as $N^{5/3}$ with the particle number N . Thus at higher density, we have the atomic helium liquid which is a Landau-Fermi liquid. As the density is increased further potential energy further decreases. At very high density the zero point energy is completely overcome, however the repulsive hard core (two body) potential comes into play. In He^3 , in the angular momentum channel $l = 0$, the repulsion is too strong. However, in $l = 1$ and $l = 2$ channel the attractive potential energy over takes the repulsion [14–16]. Thus the Lieb theory leads to 'unbounded from below' energy in the two channels (interaction in different angular momentum channel decouple). This makes the quantum mechanical Landau-Fermi liquid ground state unstable. On the other hand, Feldman theory implies that for the angular momentum channels $l = 1$ and $l = 2$, the effective running coupling constants blows up.

In the large flavour limit, the effective coupling constant in QED blows up exactly like the effective coupling constant in the Feldman theory. The Lieb treatment of the Coulomb system also suggests that in the relativistic limit for some value of the number of flavour N_f , the ground state become unstable. These similarities clearly suggest that the perturbative

vacuum state of QED is unstable. Note that in the Lieb theory for coulomb systems, we had nonrelativistic treatment followed by a relativistic system for the stability of the ground state. In condensed matter systems, the formation of cooper pairs changes the situation drastically. The liquid makes a transition to superfluid phase. The cooper pairs are essentially free and therefore, the new superfluid state does not suffer from the type of instability mentioned above. Does the instability of ground state in QED imply the existence of BCS type of ground state? The answer seems to be negative. The residue at the poles of the effective running coupling constant in QED and in the condensed matter have opposite signs. In the condensed matter system it signifies the presence of cooper pair while in the case of QED the opposite sign signifies the absence of such pairs [2,17].

The remarkable analogy of the evolution equations of the effective coupling constants in large flavour QED (Eq.(2)) and the condensed matter system above (Eq.(20)), lead to the suggestion that the perturbative ground state (vacuum state) is unstable in large flavour QED too. Landau singularity, just as in case of the condensed matter model, reflects the fact that we have constructed the perturbative theory around the "wrong" unstable ground state. In our model of QED, the elementary excitations around the perturbative vacuum or ground state are the electrons, positrons and photons and, therefore, instability of ground state or the vacuum implies that such excitations have become unstable. The new stable ground state would require modification of the particle spectrum. There are suggestions in the literature that the electric charge is totally screened by the new nonperturbative ground state [1,8,9,11,18] . The most interesting and exciting possibility is the one suggested by Witten [19]. His conjecture is that at very high energy the states carrying electric charge become very massive and are suppressed while magnetic monopoles become lighter and are excited.

Acknowledgement: I would like to thank Prof.J.Froehlich and Prof.G.M.Graf for hospitality at the Institute of Theoretical Physics, ETH- Zurich, Switzerland where part of the work was carried out.

REFERENCES

- [1] L.D.Landau and I.J.Pomeranchuk, Dokl.Akad.Nauk.,SSSR **102**,489(1955)
- [2] D.V.Shirkov, Evolution of the Bogoluibov Renormalization Group, pp 25-58 in "Quantum Field Theory: A 20th Century Profile" Ed. by A.N.Mitra, Hindustan Book Agency and Indian National Science Academy, New Delhi 2000
- [3] F.J.Dyson and A.Lenard , J.Math.Phys.(N.Y.) **8**, 423(1967); **9**, 698(1967)
- [4] E.H.Lieb *et al*, The stability of matter:From atom to stars, Selecta of E.H.Lieb, Ed. W.E.Thirring, Springer-Verlag, 1991
- [5] E.H.Lieb *et al*, Phys.Rev.Lett.**79**,1785(1997)
- [6] J.G.Conlon *et al*, Comm. Math. Phys. **116**, 417(1988)
- [7] J.Feldman, J.Magnan, V.Rivasseau, and E.Trubowitz, Euro-phys.Lett.**24**, 437 (1993); **24**, 521 (1993)
- [8] J.Kogut, E.Dagotto and A.Kocic, Phys.Rev.Lett **61**, 2416(1988);Phys.Rev.Lett **60**, 772(1988)
- [9] E.Dagotto, A.Kocic and J.Kogut, Phys.Lett. bf 232B 235(1989)
- [10] V.Miransky Nouvo Cimento **90A** 149(1985)
- [11] M.Gockeler *et al* Phys.Rev.Lett. **80** 4119(1998)
- [12] E.Lieb and H.T.Yau, Commun.Math.Pyhs bf 118 177(1988)
- [13] T.Chen, J.Froehlich, and M.Seifert, cond-mat/9508063
- [14] D.Wollhardt and P.Wolfle, The Superfluid Phases of He3, Taylor and Francis, London, 1990
- [15] K.A.Brueckner, T.Soda, P.W.Anderson, and P.Morel, Phys.Rev. **118**(1960)1442

- [16] P.W.Anderson, and P.Morel, Phys.Rev. **123** (1961)1911
- [17] A.A.Abrikosov, L.P.Gorkov, and I.E.Dzyaloshinski, Methods of Quantum Field Theory in Statistical Physics, Prentice-Hall, INC., Englewood Cliffs, New Jersey, 1963 (Chapter-I, Sec.5, Page 40)
- [18] S.Kim, J.B.Kogut, and M-P.Lombardo, Phys.Lett.B **502**, 345(2001)
- [19] E.Witten, hep-th/0212247